

**IN THE CLAIMS**

Please amend the claims, as follows, without prejudice, without admission, without surrender of subject matter, and without any intention of creating any estoppel as to equivalents:

67. (Amended) The method for breaking B-cell autotolerance in an animal to a self-protein of that animal, and inducing antibody production in the animal against the self-protein of that animal of claim 66 wherein the T-cell epitope comprises at least 10 amino acids.

68. (Amended) A method for breaking B-cell autotolerance in an animal to a self-protein of that animal, and inducing antibody production in the animal against the self-protein of that animal, comprising;

preparing different modified self-proteins, wherein:

each modified self-protein is modified, in comparison to the self-protein, by containing a substitution of a peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal,

and

the different modified self-proteins differ from each other with respect to the position of the at least one immunodominant T-cell epitope;

ascertaining which of the different modified self-proteins elicits a desired specific neutralizing effect and thereby ascertaining a desired modified self protein; and

administering to the animal, an immunologically effective amount of the desired modified self-protein, wherein:

the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein, and,

the desired modified self-protein elicits antibodies that are against the self-protein, and B-cell autotolerance to the self-protein is broken.

70. (Amended) A method for breaking B-cell autotolerance in an animal to a self-protein of that animal, and inducing antibody production in the animal against the self-protein of that animal, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier

protein or peptide containing T-cell epitopes, comprising administering to the animal, an immunologically effective amount of at least one modified self-protein, wherein:

a. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of a peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, wherein the peptide fragment corresponds to the peptide containing at least one immunodominant T-cell epitope in amino acid length;

whereby, the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

b. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of a peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, wherein the substitution is of an amphiphatic alpha helix;

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

c. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of a peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, wherein the peptide fragment corresponds to the peptide containing at least one immunodominant T-cell epitope in amino acid length and comprises at least ten amino acids;

whereby the modified self-protein elicits antibodies that are against the self-protein and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

d. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of a peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, wherein the peptide fragment corresponds to the peptide containing at least one immunodominant T-cell epitope in amino acid length and comprises at least fifteen amino acids;

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

e. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of a peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, wherein the peptide fragment corresponds to the peptide containing at least one immunodominant T-cell epitope in amino acid length and comprises at least 10 amino acids with said substitution preserving flanking regions comprising at least four amino acids on each side of the peptide fragment,

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

f. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by containing a substitution of a peptide fragment of the self-protein with a peptide containing at

least one immunodominant T-cell epitope which is foreign to the animal, wherein the peptide fragment corresponds to the peptide containing at least one immunodominant T-cell epitope in amino acid length and comprises at least 15 amino acids with said substitution preserving flanking regions comprising at least four amino acids on each side of the peptide fragment,

whereby the modified self-protein elicits antibodies that are against the self-protein earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes; and B-cell autotolerance to the self-protein is broken; or,

g. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by being detoxified by containing a substitution of a peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal,

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

h. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by being detoxified by containing a substitution of a peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, and said substitution preserving flanking regions comprising at least four amino acids on each side of the peptide fragment;

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

i. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by being detoxified by containing a substitution of a peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, wherein said peptide containing at least one immunodominant T-cell epitope comprises at least 10 amino acids and said substitution preserves flanking regions comprising at least four amino acids on each side of the peptide fragment;

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken; or,

j. the self-protein is normally autotolerated by the animal and there is normally B-cell autotolerance by the animal to the self-protein; and,

the modified self-protein is modified, in comparison to the self-protein, by being detoxified by containing a substitution of a peptide fragment of the self-protein with a peptide containing at least one immunodominant T-cell epitope which is foreign to the animal, wherein the peptide containing at least one immunodominant T-cell epitope comprises at least 15 amino acids and said substitution preserves flanking regions comprising at least four amino acids on each side of the peptide fragment,

whereby the modified self-protein elicits antibodies that are against the self-protein, earlier and in higher titres, in comparison to the self-protein conjugated to a carrier protein or peptide containing T-cell epitopes, and B-cell autotolerance to the self-protein is broken.

LD  
Cont.